



Synthesis, IR characterization and antioxidant capacity of Cu(II) complexes with amino acids and melatonin

Ljubijankić, S.^{a*}, Galijašević, S.^b, Davidović-Plavšić, B.^c, Balaban, M.^c, Ljubijankić, N.^d

^aFaculty of Health Studies, University of Bihać, Nositelja hrvatskog trolista 4, Bihać, Bosnia and Herzegovina

^bSarajevo School of Science and Technology, Sarajevo Medical School, Department of Medical Chemistry and Biochemistry, Hrasnička cesta 3b, Sarajevo, Bosnia and Herzegovina

^cFaculty of Natural Sciences and Mathematics, University of Banja Luka, Mladena Stojanovića 2, Banja Luka, B&H

^dFaculty of Science, Department of Chemistry, Zmaja od Bosne 33-35, Sarajevo, Bosnia and Herzegovina

Article info

Received: 17/02/2022

Accepted: 12/04/2022

Keywords:

Copper(II) Chloride

Cu(II) Complexes

Amino Acid

Melatonin

FT-IR

Abstract: In this paper, the reactions of anhydrous copper(II) chloride in methanolic solution with alanine, glycine, histidine, L-tryptophan, and melatonin were investigated and the resulting products were characterized by FTIR spectroscopy. Based on FTIR characterization, it was shown that these reaction conditions result in the formation of Cu(II) complexes with glycine and alanine, which are more complex structures than bis(glycinato) - and bis(alaninato) copper(II) complexes. Analysis of the FTIR spectrum of the histidine complex shows the participation of several groups in coordination at the Cu(II) center. The complex prepared with melatonin shows unusual changes in the spectral region of the amidic nitrogen bonds. The latter observation is very significant for very few known metal complexes of melatonin, while the synthesis of most of them is an experimentally demanding process with simultaneous control of several parameters. The antioxidant capacity of the synthesized complexes was examined, with the CEAC range from 121.6 to 734.4 μ M. The lowest values of antioxidant capacity were recorded for the copper complex with tryptophan, while the highest values were recorded for the copper complex with alanine. A high antioxidant capacity of the copper complex with melatonin (673.8 μ M) was also observed.

*Corresponding author:

Sead Ljubijankić

E-mail: seadlj@gmail.com

Phone: +387 037 316 701

INTRODUCTION

Due to the fact that copper is a biogenic element that performs most of its biological functions within coordination compounds with biomolecules in which, as donors of heteroatoms in the coordination sphere of the metal center, amino acids and their prosthetic groups often occur, there is a need to evaluate coordination behavior $\text{Cu}^+/\text{Cu}^{2+}$ according to individual, proteinogenic amino acids. Many reports were devoted to understanding the coordination characteristics of such complexes (Colaneri *et al.*, 2009) especially copper(II) complexes with amino acids. Copper is an essential trace element used by both eukaryotic and prokaryotic cells in anabolic and catabolic processes with fine control of the

same, given the fact that an excess of this element is associated with toxic effects. Roughly speaking, the essence of copper activity in biochemical processes is reflected in its redox behavior, especially because it can easily donate electrons to oxygen. This fact makes a certain element a potential generator of free radicals, which requires the establishment of a number of control mechanisms by the cell in order to use its reactivity to its advantage, while suppressing potential harmful effects. Therefore, the human body has established a number of biochemical processes that maintain copper homeostasis, along with homeostasis of iron, normal tissue proliferation and oxidative metabolism (Vest *et al.*, 2012). Its sudden availability in the biosphere and biologically acceptable redox potential have made

copper an essential component of oxidative metabolism (Scheiber, Dringen & Mercer, 2013). Rigorous control of copper metabolism takes place in order to prevent undesirable, copper-mediated redox processes that would ultimately create reactive oxygen species given the ease of Cu (I) / (II) conversion. Since metabolic processes involving copper have many unresolved metabolites and metabolic pathways (as well as their control), it was useful to study the coordination behavior of this biometal towards selected biomolecules under controlled conditions by the preparation and characterization of appropriate complexes. The most simple biomolecules normally present in the cell cytoplasm, i.e. the environment where copper is metabolized, are amino acids, which made them a logical starting point for studying the behavior and coordination preferences of copper in biological systems. Given the structure of proteins as important building blocks and functional components of living cells, it is certain that the potential binding of copper in one of its two oxidative states will depend on all levels of protein structure with which it interacts and on physicochemical characteristics of the cellular environment, in which the interactions take place and the product is created (Dos Santos Carvalho & Fernandes, 2019). Copper(II) complexes with amino acids have long been known, and represent a special area of interest in the coordination chemistry of this element precisely because they serve as models for the study of copper metalloproteins. In this context, the behavior of Cu(II) in aquatic systems towards histidine is particularly interesting. It has been observed that most copper binding domains of proteins are rich in this amino acid, which is consistent with the coordination preferences of copper in the +2 oxidation state towards N donor ligands. The copper(II) complex with the simplest amino acid, glycine, was prepared in 1841, by Boussingault (Delf, *et al*). It was later shown that this type of complex does not belong to conventional copper complexes precisely because of the properties of the given ligands, i.e. amino acids. Since amino acids have a larger number of potential donor atoms, the structure of the final complex certainly depends on the reaction conditions, i.e. the nature of the solvent, temperature, pH and counterion, as is the case in a living cell. Oxidative stress occurs due to an imbalance between oxidants and antioxidants in biological systems, and is caused by excessive formation of reactive oxygen species or improper functioning of the antioxidant system (Chiurchiu *et al.*, 2016). Reactive oxygen species (ROS), although possessing physiological functions, have been implicated in the pathogenesis of a number of diseases including cardiovascular and neurodegenerative diseases, muscular dysfunction, and cancer (Zuo *et al.*, 2015; He and Zuo, 2015). In addition to the many physiological effects of melatonin, melatonin has been

shown to have strong anti-oxidant properties, due to the fact that it reacts with free radicals giving other metabolites that also have a neutralizing effect on free radicals. In this way, the product of one reaction, becomes a reactant of the other in both cases consuming free radicals. Melatonin, as well as its reaction products with hydroxyl radicals, are not extremely effective in neutralizing hydroperoxyl radicals, but still suppress lipid peroxidation *in vivo*, most likely by eliminating the initial hydroxyl radicals (Galano *et al.*, 2014). In the last two decades, complexes of platinum, ruthenium, copper, cobalt and other metals have been used to modify and / or detect A β aggregation. The results of these studies suggest that metal complexes have promising potential in the treatment and diagnosis of Alzheimer's disease (Liu and Wang, 2018). Copper plays a significant role in the pathology of many neurodegenerative diseases, from Wilson and Menkes disease, through amyotrophic lateral sclerosis and Alzheimer's disease, to prion-associated diseases (Waggoner *et al*, 1999). Most of these diseases, directly or indirectly, lead to changes in copper metabolism. Changes in copper metabolism can lead to its accumulation in certain regions of the body or change its bioavailability in the cellular environment. Both are associated with increased oxidative stress and the formation of ROS, of course, due to the well-known redox behavior of copper, which the body significantly utilizes under homeostatic conditions (Waggoner *et al.*, 1999). Given the nature of melatonin as a scavenger of destructive radicals and the consequent formation of metabolites of good chelating power for Cu (Galano *et al.*, 2014) through the *scavenger cascade* of melatonin, it was interesting to examine the effect of melatonin as a potential inhibitor of Cu-protein complex species formation. Research conducted to determine the role of melatonin as a Cu(II) chelator in preventing its degrading effect on DNA (Wang *et al.*, 2019) has concluded that the copper-detoxifying and antioxidant abilities of melatonin are not based solely on scavenger abilities against hydroxyl radicals which occur in Cu(II)/Cu(I) mediated reactions, but strongly suggest the role of melatonin or its metabolites (formed by the scavenger cascade) as chelators. The final evidence lacking for unequivocal confirmation of these assumptions is the synthesis, isolation, and structural characterization of such complex types of melatonin, its metabolites, and copper.

EXPERIMENTAL

Materials and methods

All chemicals used were commercially available with analytical grade of purity and used without further purification. Infrared spectra were recorded as KBr pellets on a Perkin Elmer spectrum BX FTIR System in

region 4000 – 400 cm^{-1} , resolution 2cm^{-1} , 8 scans. The antioxidant activity was determined using the e-BioQuChem (e-BCQ) lab device, following the manufacturer's instruction manual. The results are expressed in μC or in the charge of the electrons released by the antioxidant to neutralize free radicals. This device makes it possible to distinguish fast-acting antioxidants such as ascorbic acid and coenzyme Q10 from co-producing antioxidants such as polyphenols, astaxanthin or α -lipoic acid. Fast-acting antioxidants are the first to be oxidized and are considered to be more potent antioxidants than general antioxidants even though they are present in lower concentrations. Total antioxidant capacity (QT) is obtained as the sum of the antioxidant capacity of the fast-acting (Q1) and co-producing antioxidants (Q2). The results of antioxidant capacity were also expressed in ascorbic acid equivalents (CEAC), for which it was necessary to construct a calibration curve. For this purpose, a dilution series of a solution of ascorbic acid standard in 0.1 M PBS buffer (pH = 7) was prepared in the concentration range 0 - 1000 μM .

Synthesis of complexes

All complexes were synthesized by heating the reaction mixture to 65°C on a magnetic stirrer, under reflux for 5 to 9 hours. Immediately before the syntheses, anhydrous copper(II) chloride (CuCl_2) was prepared by drying copper(II) chloride dihydrate ($\text{CuCl}_2 \times 2\text{H}_2\text{O}$) at 120°C for 2 hours. This drying gave anhydrous CuCl_2 , which was observed by changing the color from blue, copper(II) chloride dihydrate to brown colored, anhydrous copper(II) chloride. For the synthesis of each complex, solutions of copper(II) chloride in methanol were prepared by dissolving 268.9 mg (0.002 mol) of anhydrous CuCl_2 in 40 mL of methanol preheated to 50°C .

Synthesis of Cu(II) complex with alanine

In 40 mL of methanol previously heated to 50°C , 356.36 mg (0.004 mol) of alanine were dissolved and a previously prepared solution of copper(II) chloride was added to the solution. After mixing, the solution was stirred a magnetic stirrer at 65°C under reflux for about 5 hours. Immediately after mixing the solution, the reaction mixture showed light green color. After three hours of heating, the color of the mixture turned dark green and did not change until the end of the synthesis. After cooling, the reaction mixture was filtered and the crystals of the complex were washed with water and ether. Finally, the complex was recrystallized from dimethylsulfoxide/water and dried in a vacuum desiccator.

Synthesis of Cu(II) complex with glycine

The amount of 303.30 mg (0.004 mol) of glycine was dissolved in 40 mL of methanol preheated to 50°C , and the previously prepared copper(II) chloride solution was added to the solution. The reaction mixture was heated with a magnetic stirrer at 65°C under reflux for about 5 hours. Immediately after mixing the solution, the reaction mixture had a light blue color which eventually turned a slightly darker shade of blue. The mixture was cooled and left at room temperature for 24 hours. The blue crystals of the resulting complex were then filtered and washed with water and ether, recrystallized from dimethylsulfoxide/water and dried in a vacuum desiccator.

Synthesis of Cu(II) complex with histidine

To 620.62 mg (0.004 mol) of histidine was added 40 mL of methanol preheated to 50°C . To the resulting solution was then added a previously prepared solution of copper(II) chloride. During the reaction, the color of the reaction mixture changed from light green-blue at the beginning to dark purple-blue at the end of the synthesis. After cooling and filtering the mixture, the crystals of the complex were washed with water and ether, recrystallized from dimethylsulfoxide/water and dried in a vacuum desiccator.

Synthesis of Cu(II) complex with L-tryptophan

For synthesis, 816.92 mg (0.004 mol) of L-tryptophan were weighed and dissolved in 50 mL of methanol heated to about 50°C . CuCl_2 solution was added dropwise to the resulting solution with continuous stirring. The mixture was heated at 65°C on a magnetic stirrer under reflux for 5 hours. After the end of the synthesis, the copper(II) complex with L-tryptophan was deposited in the form of a powdery substance of dark brown-blue color. The reaction mixture was filtered and the complex washed with hot water to eliminate chlorides, recrystallized from dimethylsulfoxide/water and dried on filter paper in a vacuum desiccator.

Synthesis of Cu(II) complex with melatonin

For the synthesis, 929.11 mg (0.004 mol) of melatonin was dissolved in 40 mL of methanol heated to 50°C . A previously prepared solution of copper(II) chloride was added to the resulting solution. Immediately after mixing, the reaction mixture turned greenish brown. The mixture was heated to 65°C while stirring with a magnetic stirrer under reflux for 9 hours. During the synthesis, the color of the mixture changed, to eventually become dark brown, which did not change until the end of the synthesis. After cooling and filtering the mixture, the crystals of the complex were washed with water and petroleum ether, recrystallized from dimethylsulfoxide/water and dried in a vacuum desiccator.

RESULTS AND DISCUSSION

All complexes were synthesized by the reaction of methanolic solutions of CuCl_2 and the corresponding amino acids (alanine, glycine, histidine and tryptophan), and melatonin, at 65°C under reflux for 5 to 9 hours.

FTIR spectra of amino acids and melatonin

The FTIR spectra of all amino acids show vibrations typical of the carboxyl group, whose displacements in different amino acids are attributed to its protonated or deprotonated form. The asymmetric stretching of the carboxyl group in the given amino acids occurs at about 1600 cm^{-1} and is extremely important for determining the performance and type of coordination in Cu-complexes. The area above 3000 cm^{-1} is not of special importance for free amino acids, but it becomes more important for their Cu-complexes. Vibrations in the amino acid and melatonin spectra in the range of 2000 to 2800 cm^{-1} are not of particular importance in determining subsequent coordination at the metal center. Extremely significant vibrations in the spectra of amino acids, but also the metal complexes is that attributed to the amino group in either protonated or deprotonated form. These bands are found in alanine at about 1590 , 1520 and 1230 cm^{-1} . Indispensable are the vibrations of the hydrocarbon skeleton that occur at about 1000 cm^{-1} depending on the type of amino acid, as well as those attributed to the aromatic fragment of the molecule in the case of histidine, tryptophan and melatonin. Vibrations of C-H bonds for the aliphatic part of the molecule occur below 3000 to 2900 cm^{-1} , while for the aromatic part of the molecule they occur above 3000 cm^{-1} to 3100 cm^{-1} and are clearly visible in the spectra of histidine, tryptophane and melatonin. The expected deviations in spectral characteristics in relation to amino acids show melatonin as a typical amine where this functionality is shown at about 3280 cm^{-1} and at 1490 cm^{-1} , while the indole functionality is found at 3305 cm^{-1} and 1270 cm^{-1} , respectively. Other bands in the ligand spectra have a more specific character.

FTIR spectra of Cu(II) complexes

Copper spectra with amino acids and melatonin show typical low-band bands attributed to metal-ligand vibrations and are of great importance for determining the coordination type and potential geometry of the complex. As copper(II) chloride was used as the starting substance, the retention of chloride in the coordination sphere in the final complex is not excluded. The previously mentioned typical vibrations of amino acids and melatonin, which now play the role of ligands, show typical shifts that indicate a model of coordination at the Cu(II) center. A typical phenomenon after coordination is the abolition of individual bands or the abolition of degenerate vibrational modes. A typical, spectrally noticeable change shown by a coordinated carboxyl

group is the appearance of a band corresponding to the C=O bond as well as a band corresponding to the C-O bond, which indicates the abolition of resonance in the uncoordinated carboxyl group, i.e. its participation in coordination to the metal center. An additional change in relation to the spectra of free ligands is often the deprotonation of the amino group due to its coordination, which is observed by the abolition of bands typical of the $-\text{NH}_3$ group and the appearance of bands typical of the $-\text{NH}_2$ group. Skeletal vibrations do not suffer significant displacements or their bands are canceled if a certain vibration mode becomes impossible after coordination with the metal center, which is also important data. Other bands in the spectra of the complex have a more specific character and are described in detail for each individual complex.


FTIR characterization of Cu - alanine complex

It is known that the reaction of copper(II) acetate, as a salt of a weak organic acid, with alanine in adequate solvents and with good pH control creates bis(alaninato) copper(II) complex which occurs in two geometric isomers, ie *trans*-bis alaninate) copper(II) monohydrate and *cis*-bis(alaninato) copper(II) monohydrate. The difference between these two isomers can be easily determined by IR spectroscopy, with vibrations in the spectral range of 750 to 200 cm^{-1} bei considered relevant. The *trans* isomer generally shows fewer bands in said region as well as *trans*-bis(glycinato) copper(II). In essence, metal-ligand vibrations may be sufficient to perform this distinction (Herlinger, Wenholt, & Long, 1970) where $\nu(\text{Cu-N})$ vibration occurs as a weak band at 484 cm^{-1} in the case of the *trans* isomer, while the *cis* isomer shows $\nu(\text{Cu-N})$ at 483 cm^{-1} and $\nu(\text{Cu-N})$ at 411 cm^{-1} . In the case of the prepared Cu complex with alanine $\nu(\text{Cu-N})$ vibration is observed at 472 cm^{-1} , which is not correlated with typical bis(alaninato) copper(II) complexes. A wide band in the spectrum of the complex at about 3429 cm^{-1} as well as an intense band at 1637 cm^{-1} are used to detect crystal bound water, the former being attributed to vibrations of O-H bond stretching water molecules and the latter to their symmetric deformation (Nakamoto, 2009). Bands in the range $3297 - 3140\text{ cm}^{-1}$ were attributed to the (N-H) vibrations of the coordinated amino group. These bands are more visible in the complex in relation to the vibrations of the same bond of the protonated amino group in the ligand. The deformation vibration of this group is in the vicinity of the band for C=O bonding and deformation of water molecules, and has a peak at 1598 cm^{-1} (Han, 2010). Vibration of the protonated amino group, i.e. $\nu_r(\text{NH}_3^+)$, is not present in the spectrum of the complex, and has been replaced by "rocking" and "wagging" vibrations of the coordinated amino group at 1150 cm^{-1} and 1022 cm^{-1} (Nakamoto, 2009) the intense band at 666 cm^{-1} also attributes the "rocking" vibration

to this group. The band at 1578 cm^{-1} corresponds to the vibration of the $\nu(\text{C}=\text{O})$ coordinated carboxyl group, while the $\nu(\text{C}-\text{O})$ vibration occurs at 1396 cm^{-1} . Resonance of the carboxyl group is reflected in the alanine spectrum by the $\nu_{as}(\text{COO}^-)$ vibration at 1624 cm^{-1} . This band is not present in the spectrum of the complex, which confirms coordination through the carboxyl group. The bands corresponding to the in-plane or "out-of-plane" vibrations of the carboxyl group at 850 cm^{-1} and 650 cm^{-1} , as well as the deformation at 772 cm^{-1} disappear in the spectrum of the complex and are replaced by "rocking" or wagging by vibrations of a coordinated carboxyl group at 530 cm^{-1} and 624 cm^{-1} as much less intense bands (Herlinger, Wenhold and Long, 1970). Bands whose positions and intensities have little or no change in the spectrum of alanine and the respective complex originate mostly from C-H vibrations and deformations of alkyl groups. Namely, the bands at 2927 cm^{-1} and 2856 cm^{-1} in the spectrum of the complex were attributed to the (C-H) vibrations of the methyl and (CH) groups (Garcia *et al.*, 2008). The band at 1113 cm^{-1} attributed to $\nu_{as}(\text{C}-\text{C}) + \rho(\text{NH}_3^+)$, which is in a complex of lower intensity and occurs 1116 cm^{-1} , reflects a higher proportion of C-C vibration. Bands of symmetrical and asymmetric deformation of the methyl group in the prepared complex 1366 cm^{-1} and 1460 cm^{-1} occur, almost unchanged, relative to the ligand. Bands corresponding to free vibrations of the C-C-N skeleton do not appear in the spectrum of the complex, which is a consequence of coordination through N atoms (Nakamoto, 2009).

Table 1: Characteristic vibrations of Cu(II) alanine complex

Characteristic vibrations of Cu-glycine	Wave number of vibration, ν (cm^{-1})
$\nu(\text{O}-\text{H})\text{H}_2\text{O}$	3464 b
$\delta_s(\text{H}_2\text{O})$	1640 s
$\nu(\text{N}-\text{H})\text{NH}_2$	3356 – 3250 m, b
$\nu(\text{C}-\text{H})$	2984 – 2924 w – m
$\delta(\text{NH}_2)$	1602 s
$\nu(\text{C}=\text{O})$	1578 s
$\delta_s(\text{NH}_3^+)$	–
$\delta_s(\text{CH}_2)$	1444 s
$\nu(\text{C}-\text{O})$	1408 s
$\rho_w(\text{CH}_2)$	1338 m
$\nu_r(\text{NH}_3^+)$	–
$\rho_r(\text{NH}_2)$	1101 m-s
$\rho_w(\text{NH}_2)$	1060 m
$\nu_{as}(\text{C}-\text{C}-\text{N})$	–
$\rho_r(\text{CH}_2)$	916 w
$\nu_s(\text{C}-\text{C}-\text{N})$	–
$\rho_w(\text{COO}^-)$	–
$\delta(\text{C}=\text{O})$	742 w
$\pi(\text{C}=\text{O})$	608 m
$\nu(\text{Cu}-\text{N})$	486 m

**Figure 1:** Comparative FT-IR spectra of alanine and Cu- alanine complex

FTIR characterization of Cu - glycine complex

It is known that the reaction of copper(II) acetate, as a salt of a weak organic acid, with glycine in adequate solvents and with good pH control creates bis (glycinato) copper(II) complex which occurs in two geometric isomers, ie trans-bis (glycinato) copper(II) dihydrate and cis-bis (glycinato) copper(II) monohydrate. IR spectroscopy can easily determine the true difference between these two isomers, with vibrations in the spectral range of 750 to 200 cm^{-1} being considered relevant. Having greater symmetry, the trans isomer generally shows fewer bands in said region. In essence, metal-ligand vibrations may be sufficient to perform this distinction (Herlinger, Wenholt and Long, 1970) where $\nu(\text{Cu-N})$ vibration occurs as a weak band at 477 cm^{-1} in the case of the trans isomer, while the cis isomer shows $\nu_{\text{as}}(\text{Cu-N})$ at 471 cm^{-1} and $\nu_{\text{s}}(\text{Cu-N})$ at 454 cm^{-1} . In the case of the prepared Cu complex with glycine $\nu(\text{Cu-N})$ vibration is observed at 486 cm^{-1} , which is not correlated with typical bis (glycinato) copper(II) complexes. A wide band in the spectrum of the complex at about 3464 cm^{-1} with an intense band at 1640 cm^{-1} is used to prove crystal bound water where the former is attributed to vibrations of stretching O-H bonds of water molecules and the latter to their symmetric deformation (Nakamoto, 2009). The bands at 3169 cm^{-1} in the glycine spectrum attributed to the (N-H) protonated amino group were replaced by bands in the range of 3356 - 3250 cm^{-1} , which are attributed to the (N-H) vibrations of the coordinated amino group. These vibrations for *trans*-bis (glycinato) copper(II) dihydrate occur in the range of 3320 - 3260 cm^{-1} , which does not correlate with the bands of the synthesized Cu-glycine complex, which is therefore more complex than simple bis (glycinato) copper (II) complex. The band at 1578 cm^{-1} corresponds to the $\nu(\text{C}=\text{O})$ coordinated carboxyl group, while the $\nu(\text{C}-\text{O})$ vibration occurs at 1408 cm^{-1} , which confirms the coordination of the carboxyl group, as well as the resonance that occurs in the glycine spectrum by vibration at 1612 cm^{-1} which is not present in the spectrum of the complex. The bands corresponding to the "rocking" or "wagging" vibrations of the carboxyl group at 508 cm^{-1} and 698 cm^{-1} , as well as the deformation at 610 cm^{-1} , disappear in the spectrum of the complex and are replaced by much less intense vibration bands of the coordinated carboxyl group, at 742 cm^{-1} and 608 cm^{-1} (Inomata *et al.*, 1988). Bands whose positions and intensities have little or no change in the spectrum of glycine and its complex originate mostly from C - H vibrations and deformations of the CH_2 group.

Namely, bands in the range of 3000 to 2900 cm^{-1} appear in both the ligand and the complex, and are attributed to $\nu(\text{C-H})$ vibrations. There is a band "wagging" vibrations of the CH_2 group at about 1337 cm^{-1} practically unchanged position in both substances, but much lower intensity in the complex. This change in intensity can be related to electronic changes due to coordination on the charged metal center. The same can be concluded for the "wagging" vibration of the CH_2 group, which was shifted from 910 cm^{-1} in the ligand to 916 cm^{-1} in the complex, and significantly reduced in intensity. It is clear that coordination limits the ease of development of certain types of vibrations (especially deformation vibrations and torsional vibrations), which is combined with electrostatic interactions in the crystal lattice of a complex joint and may justify small changes in band positions reflecting vibrations. such as the CH_2 group of glycine (Nakamoto, 2009).

Table 2. Characteristic vibrations of Cu(II)-glycine complex

Characteristic vibrations of Cu-alanine	Wave number of vibration, ν (cm^{-1})
$\nu(\text{O-H})\text{H}_2\text{O}$	3429 b
$\delta_{\text{s}}(\text{H}_2\text{O})$	1637 s
$\nu(\text{N-H})\text{NH}_2$	3297 – 3140 m, b
$\nu(\text{C-H})$	2927, 2856 w – m
$\delta(\text{NH}_2)$	1598 m
$\nu(\text{C=O})$	1578 s
$\delta_{\text{as}}(\text{CH}_3)$	1460 s
$\delta_{\text{s}}(\text{NH}_3^+)$	–
$\delta_{\text{s}}(\text{CH}_3)$	1366 s
$\nu(\text{C-O})$	1396 s
$\nu_{\text{as}}(\text{C-C})$	1116 m
$\nu_{\text{r}}(\text{NH}_3^+)$	–
$\rho_{\text{r}}(\text{NH}_2)$	1150 w
$\rho_{\text{w}}(\text{NH}_2)$	1022 w
$\delta(\text{COO}^-)$	784, 722 w
$\rho_{\text{r}}(\text{NH}_2)$	666 m
$\rho_{\text{w}}(\text{COO}^-)$	624 m
$\rho_{\text{r}}(\text{COO}^-)$	530 w
$\nu(\text{Cu-N})$	472 m




Figure 2: Comparative FT-IR spectra of glycine and Cu- glycine complex

FTIR characterization of Cu - histidine complex

In terms of coordination chemistry, histidine has been described as a rather complex ligand. Due to the large number of adequately positioned donor atoms, histidine as a ligand can build various complicated complexes with metals that prefer N donor ligands, such as Cu(II). Kruck and Sarkar (1973) described at least six different coordination forms in the Cu(II) histidine system where histidine always occurs as an O, N donor. The N donor atom may come from the amino group of the aliphatic chain or the N atom of the imidazole ring carrying a free electron pair. In the spectrum of the complex at 418 cm^{-1} , the band attributed by Drodźewski and Kordon (2000) to $\nu(\text{Cu-N})$ occurs when the N atom originates from the amino (NH_2) group and the assumption was confirmed by isotopic derivatization. This, however, does not exclude the occurrence of histidine as a Nimidazole donor, but only confirms that histidine also acts as an N (NH_2) donor in the prepared complex. Coordination to the Cu(II) center via the amino group is supported by changes in other bands, fully or partially attributed to the vibrations of this group in free histidine. Namely, the band attributed to the "scissoring" vibration of the amino group at about 1632 cm^{-1} was moved to 1625 cm^{-1} , while the band attributed to $\rho\omega(\text{NH}_2)$ at 925 cm^{-1} in the spectrum of the complex disappears. The amino group vibration "rocking" band combined with other histidine vibrations is present at 827 cm^{-1} , but expanded and drastically less intense, which also suggests coordination across the amino group (Nakamoto, 2009). The intense band attributed to the combined deformation of the N-H and C-H bonds of the imidazole fragment of the

molecule in the histidine spectrum disappears in the spectrum of the complex, which may indicate the participation of the imidazole fragment in coordination with the Cu(II) center. More definite evidence would be found in Cu-Nimidaz. which occurs at about 280 cm^{-1} (Drodźewski and Kordon, 2000) which is beyond the scope of the performed recordings. The band carrying the highest proportion of carboxyl group vibrations, with a smaller proportion of other histidine vibration modes, was shifted from 1796 cm^{-1} from the ligand spectrum to 1740 cm^{-1} in the complex spectrum (Kruck and Sarkar, 1973), the most significant indicator of histidine coordination over carboxyl groups (Dunbar *et al.*, 2018). It should be noted that the Cu(II) complex that two histidine molecules can have multiple forms, one of which contains Cu(II) in the tetragonal distortion of O, N, N donor histidine molecules. The intense band at 625 cm^{-1} attributed to the out-of-plane bending of the N-H bond of the imidazole ring (combined with other vibrations) occurs virtually unchanged in the spectra of both ligands and complexes, which is expected given that this bond remains preserved in complex. Significant is the behavior of the band at 1453 cm^{-1} , which is partly attributed to the deformation of the CH_2 group, and in the complex occurs at approximately 1400 cm^{-1} . This decline is atypical for the group not directly involved in coordination, however the band of this vibration strongly depends on the spatial arrangement of parts of the histidine molecule, so it occurs in different conformations ranging from 1477 to 1424 cm^{-1} (Kumar *et al.*, 2010). This band shift in the spectrum of the complex relative to the ligand can be interpreted as

additional evidence of histidine coordination at the Cu(II) center, given that the coordination process almost always causes changes in the conformation of more complex ligands (Nakamoto, 2009). the position of the "scissoring" band of the vibration of the CH₂ group of histidine.

Table 3. Characteristic vibrations of Cu- histidine complex

Characteristic vibrations of Cu-histidine	Wave number of vibration, ν (cm ⁻¹)
$\nu(C=O) + \delta(O-H) + \delta(C-H)$	1740 m
$\delta_s(NH_2)$	1625 s
$\delta_s(CH_2)$	1398 s
$\delta(C-H) + \delta(N-H)$	-
$\nu(C-NH_2) + \delta(C-H)$	1084 m
$\rho_w(NH_2)$	-
$\rho_r(NH_2)$	827 m, b
$\gamma(N_{imidaz.-H})$	625 s
$\nu(Cu-N)$	418 s

FTIR characterization of Cu - tryptophan complex

The spectrum of the prepared complex shows significant differences in the bands attributed to vibrations in which nitrogen atoms participate, which leads to the first conclusion about its coordination engagement in the complex. The vibration-corresponding bands (NH₃⁺) of the group completely disappear in the spectrum of the complex and are replaced by vibrations characteristic of the NH₂ group. Thus, wild very intense bands appear at 3338 cm⁻¹ and 3254 cm⁻¹, which correspond to asymmetric and symmetric stretching of the amino group, while at 1627 cm⁻¹ there is an intense band of strain deformation in the mentioned group. Vibrations involving the indole nitrogen atom have no significant change in position or intensity. In the vibrations of the nitrogen atom that is directly related to the aliphatic carbon atom, a small shift towards higher wave numbers was observed. Vibration at 3420 cm⁻¹ indicates that the nitrogen atom in the indole ring is not deprotonated, and therefore probably not coordinated (Wagner & Baran, 2004). Equally significant changes are observed in the vibration bands of the carboxyl group where symmetrical and asymmetric stretching is not present in the molecule of the complex, thus pointing to the violation of the symmetry of the group and the abolition of resonance between bonds to a single C-O bond and a double C=O bond.

Proof of this is the very intense band at 1562 cm⁻¹ corresponding to the carbonyl bond from the carboxyl group while at 1382 cm⁻¹ the rest of the group occurs in the form of C-O vibration. The intense deformation vibration of this group was shifted toward lower wave numbers at 740 cm⁻¹ (Wagner & Baran, 2004). Vibrations corresponding to the hydrocarbon skeleton do not change significantly in position or are eliminated due to differences in the symmetry of the coordinated and free molecules of tryptophan. Metal ligand vibrations are the most convenient form of confirmation of coordination both in terms of the nature and identity of coordinated atoms. Thus, in the spectrum of the complex, typical stretching vibrations of the Cu-N bond occur, which copper forms with the deprotonated amino group of tryptophan, at 475 cm⁻¹ as a medium intensity band. Vibrations of Cu-O usually fall below 400 cm⁻¹ (about 350 - 310 cm⁻¹) so that they are not visible in the given spectrum, but their existence is indicated by changes in the carboxyl group in terms of resonance cancellation due to binding of one oxygen atom from the given group to the Cu center and the accompanying formation of the C=O bond (Nakamoto, 2009; Wagner & Baran, 2004).

Table 4: Characteristic vibrations of Cu- tryptophan complex

Characteristic vibrations of Cu-tryptophan	Wave number of vibration, ν (cm ⁻¹)
$\nu(N-H)_{indole}$	3420 vs
$\nu_{as}(NH_2)$	3338 vs
$\nu_s(NH_2)$	3254 vs
$\nu(NH_3^+) + \nu(C-H)$	-
$\nu(C-H)$	2895 w
$\delta_{as}(NH_3^+)$	-
$\delta(NH_2)$	1627 vs
$\nu_{as}(COO^-)$	-
$\nu(C=O)$	1562 vs
$\delta_s(NH_3^+)$	-
$\delta(CH_2)$	1458 s
$\nu_s(COO^-)$	-
$\nu(C-O)$	1382 m
$\tau(CH_2)$	1228 m
$\nu(C-N)_{aliph.}$	1100 m
$\delta(COO^-)$	740 s
$\nu(Cu-N)$	475 m




Figure 3: Comparative FT-IR spectra of histidine and Cu-histidine complex




Figure 4: Comparative FT-IR spectra of tryptophan and Cu-tryptophan complex

FTIR characterization of Cu-melatonin complex

The spectrum of the Cu-melatonin complex shows a band at 3308 cm^{-1} which corresponds to the vibration of the N-H bond of the indole. The appearance of this band, with small changes in symmetric and asymmetric stretching of the C - N bond within the indole ring, leads to the conclusion that this group is intact in the final complex and rejects the possibility of melatonin coordination at the Cu(II) center via N indole atoms (Shimazaki *et al.*, 2009). The band carrying the largest share of C=O vibration of the amide group occurs 1632 cm^{-1} , is lower in intensity than in the melatonin spectrum, but certainly means that the oxygen atom does

not participate in the coordination sphere of the Cu(II) center. This is supported by the sensitivity of this vibration to coordination via the amide oxygen atom, where a shift to higher wave numbers of up to 20 cm^{-1} is expected (Ashok, *et al.*, 2006). In contrast to the aforementioned groups, significant changes occur in bands involving vibrations of the amide N atom in the spectra of the melatonin and Cu-melatonin complexes. To evaluate coordination across an amide N atom, $\nu_{\text{as}}(\text{C-N})$ amide. + $\delta(\text{C-N-H})$ amide, is the most significant band corresponding to the vibration of the amide group and occurs in the spectrum of melatonin at 1490 cm^{-1} .

This band is lost in the spectrum of the complex from this spectral region. The band at 954 cm^{-1} in the melatonin spectrum, corresponding to the $\nu(\text{C-N})$ amide vibration also disappears in the spectrum of the complex. Vibration $\gamma(\text{N-H})$ amide + $\tau(\text{HCCO})$ amide, which carries a higher proportion of torsional vibrations, changes position and intensity in the spectrum of the complex. These observations suggest the possibility of coordination of melatonin at the Cu(II) center via the N atom of the amide group, which is a common tendency of amide ligands (Marti *et al.*, 2012) in the context of Cu(II) coordination chemistry. In the spectrum of the complex, the I-band appears at 480 cm^{-1} , which is the area of the IR spectrum where the stretching of the Cu-N bond is expected (Nakamoto, 2009). An extremely interesting phenomenon is the band at 1714 cm^{-1} which is not a common vibrational region of the carbonyl group attached to the N atom as is the case in amides. Ji *et al.* (2020) provided an interesting overview of the theoretically predicted spectra for simple amide (*N*-methylacetamide) for each individual resonant form. A resonant form that represents said molecule as acetimic acid (where the H atom of the amide group is attached to oxygen while carbon and oxygen form an imine bond

(C=N)) shows a stretching vibration of the C=N bond at about 1700 cm^{-1} . Coordination at the metal center significantly affects the balance of individual resonant forms, in the direction that favors better coordination with the metal center through the heteroatom that participates in the resonance (Nakamoto, 2009). It can be assumed that the Cu(II) center can coordinate to the N atom of the amide part of the melatonin molecule to induce the formation of (C=N) bonds by stabilizing such a resonant form, which would be manifested by a band at 1714 cm^{-1} . Many other experimental data are needed to finally confirm this assumption. Bands of groups that do not participate in coordination occur virtually unchanged in the spectrum of the complex relative to the spectrum of the ligand. Asymmetric and symmetric stretching of the C-H group of the aliphatic part of the molecule occurs in the complex at 2926 cm^{-1} and 2866 cm^{-1} , while the "rocking" vibration band of the mentioned bond is also practically unchanged in the complex at 1045 cm^{-1} . The band at 1555 cm^{-1} corresponding to the asymmetric stretching of the C=C bond of the indole ring at 2C - 3C is of lower intensity but of the same position, at 1556 cm^{-1} in the spectrum of the complex (Singh *et al.*, 2014).

Table 5: Characteristic vibrations of Cu - melatonin

Characteristic vibrations of Cu-melatonin complex	Wave number of vibration, ν (cm^{-1})
$\nu(\text{N-H})_{\text{indole}}$	3308 s
$\nu(\text{N-H})_{\text{amide}}$	-
$\nu_{\text{as}}(\text{C-H})_{\text{aliph.}}$	2926 m
$\nu_{\text{s}}(\text{C-H})_{\text{aliph.}}$	2866 m
$\nu(\text{C=O}) + \delta(\text{C-N-H})_{\text{amide}}$	1632 s
$\nu_{\text{as}}(\text{C=C})_{2-3\text{C-indole}}$	1556 m
$\nu_{\text{as}}(\text{C-N})_{\text{amide}} + \delta(\text{C-N-H})_{\text{amide}}$	-
$\nu_{\text{s}}(\text{C-N})_{\text{indole}}$	1264 w
$\nu_{\text{as}}(\text{C-N})_{\text{indole}}$	1206 s
$\nu(\text{C-O})_{\text{methoxy}}$	1174 m
$\rho_{\text{r}}(\text{C-H})_{\text{aliph.}}$	1045 ms
$\nu(\text{C-N})_{\text{amide}}$	-
$\gamma(\text{C-H})_{\text{indole}} + \tau(\text{HCCC})_{2-3\text{C-indole}}$	803 m
$\tau(\text{HCCO})_{\text{amide}}$	604 m
$\gamma(\text{N-H})_{\text{amide}} + \tau(\text{HCCO})_{\text{amide}}$	538 w
$\nu(\text{Cu-N})$	480 w




Figure 5: Comparative FT IR spectra of melatonin and Cu- melatonin complex

Determination of antioxidant capacity

By examining the antioxidant capacity of the synthesized copper complexes, CEAC values in the range of 121.6 - 734.4 μM were obtained. The lowest values of antioxidant capacity were recorded in the copper complex with tryptophan, while the highest values were recorded in the copper complex with alanine. A high antioxidant capacity of the copper complex with

melatonin (673.8 μM) was also observed. Melatonin is known to be a good scavenger of hydroxyl radicals, superoxide anion radicals and nitric oxide and is one of the most effective lipophilic antioxidants (Simunkova *et al.*, 2019).





Figure 6: Calibration curve with standard ascorbic acid solution

Ascorbic acid is the strongest natural antioxidant and is commonly used in determining antioxidant capacity. To present the results as ascorbic acid equivalents (CEAC),

a calibration curve was constructed as shown in Figure 6. The values of Q1 and Q2, as well as QT of the analyzed complexes are shown in Table 6.

Table 6: Antioxidant capacity of synthesized complexes expressed in μC

Complex	Q1 (μC)	Q2 (μC)	QT (μC)
Cu-Gly	3.6	20.1	23.7
Cu-Ala	24.3	50.2	74.5
Cu-His	3.9	47.2	51.1
Cu-Mel	36.2	32.6	68.8
Cu-Trp	3.9	13	16.9

**Figure 7:** Total antioxidant capacity (CEAC) of synthesized complexes

CONCLUSION

Five complex compounds of copper(II) were prepared following an identical procedure. Four amino acids (glycine, alanine, tryptophan and histidine) and melatonin as a bioactive tryptophan derivative were used as ligands. It was found that this preparative approach in the case of alanine and glycine does not result in the isolation of well-known *cis* / *trans*-bis (glycinato) copper(II) or *cis* / *trans*-bis (alaninato) copper(II) complexes implying the importance of starting copper(II) salt as well as pH control of the solution, given that said complexes are obtained using copper(II) acetate in approximately neutral aqueous solutions. FTIR characterization of the isolated complexes with glycine and alanine showed deviations from the IR spectral characteristics of the isomeric amino acid carboxylates of copper(II), especially in the region from 800 to 400 cm^{-1} where isomer-specific vibrations are common.

Analysis of the FTIR spectra of prepared complexes of these amino acids provides data on the coordination of both oxygen atoms from the carboxyl group and nitrogen atoms from the amino group of the same, but does not dispute the participation of chlorides and water molecules in the coordination sphere of Cu(II). The histidine complex also shows the participation of oxygen and nitrogen atoms from the aliphatic part of this amino acid, but based on changes in skeletal ring vibrations, potential coordination of nitrogen atoms from the imidazole residue of this amino acid is assumed, which could be concluded based on Cu(II) center coordination preference. In the case of the melatonin complexes, the FTIR spectrum disputes coordination via oxygen atoms either from the amide group of the aliphatic moiety or from the methoxyl substituent on the indole ring, the potential deprotonation of the indole ring and coordination via the nitrogen atom from the same is also challenged. In contrast, the FTIR spectrum of the

complex clearly indicates the coordination of nitrogen atoms from the amide group as well as the potential stabilization of one of the resonant forms of this group based on theoretical considerations. It is important to note that the assumed structures contain coordinated chlorides based on theoretical knowledge and practical data on the behavior of such systems, and that the final judgment on their coordination participation is made by considering the spectral range from 400 to 200 cm⁻¹ not examined in this work. Finally, it can be said that this simple preparative approach results in complex compounds of amino acids and their selected derivatives with copper(II) chloride that are different from the literature described, relatively simple complexes and that additional characterization is needed to establish the final structure of the complex and possible conclusions about mechanistic processes that are ongoing in the given reaction system. The obtained results of the antioxidant capacity of the complexes indicate a possible good antioxidant effect of the copper complex with melatonin and alanine, which indicates the need of further research in terms of the use of other methods to determine antioxidant capacity (Romero-Canelon and Sadler, 2013; Van Rijt and Sadler, 2009; Leung et al., 2015).

REFERENCES

- Ashok, M., Ravinder, V., Prasad, A. V. S. S. (2006) Synthesis, spectral characterization and catalytic applications of Ru(II) complexes with amide ligands. *Transition Metal Chemistry*, 32(1): 23–30. DOI:10.1007/s11243-006-0123-0.
- Colaneri M. J., Vitali J., and Peisach J. (2009). Aspects of Structure and Bonding in Copper–Amino Acid Complexes Revealed by Single-Crystal EPR/ENDOR Spectroscopy and Density Functional Calculations, *The Journal of Physical Chemistry*, 113 (19), 5700–5709 DOI: 10.1021/jp811249s
- Chiurchiù, V., Orlacchio, A., & Maccarrone, M. (2016). Is modulation of oxidative stress an answer? The state of the art of redox therapeutic actions in neurodegenerative diseases. *Oxidative medicine and cellular longevity*, 2016. DOI: [10.1155/2016/7909380](https://doi.org/10.1155/2016/7909380)
- Delf, B. W., Gillard, R. D., & O'Brien, P. (1979). The isomers of α -amino-acids with copper(II). Part 5. The cis and trans isomers of bis(glycinato)copper(II), and their novel thermal isomerization. *J. Chem. Soc., Dalton Trans.*, (8), 1301–1305. DOI:10.1039/dt9790001301
- Dos Santos Carvalho, C., Fernandes, M. N. (2019) Effects of copper toxicity at different pH and temperatures on the in vitro enzyme activity in blood and liver of fish, *Prochilodus lineatus*. *Molecular Biology Reports*, 46(5): 4933–4942 DOI:10.1007/s11033-019-04944-y.
- Drożdżewski, P., Kordon, E. (2000) Isotopic studies of the metal–ligand vibrations in histamine complexes with Copper(II). *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 56(7): 1299–1304. DOI:10.1016/s1386-1425(99)00232-2.
- Dunbar, R. C., Martens, J., Berden, G., Oomens, J. (2018) Transition metal(II) complexes of histidine-containing tripeptides: Structures, and infrared spectroscopy by IRMPD. *International Journal of Mass Spectrometry*, 429: 198–205. DOI:10.1016/j.ijms.2017.10.004.
- Galano, A., Medina, M. E., Tan, D. X., Reiter, R. J. (2014) Melatonin and its metabolites as copper chelating agents and their role in inhibiting oxidative stress: a physicochemical analysis. *Journal of Pineal Research*, 58(1): 107–116. DOI:10.1111/jpi.12196.
- Galano, A., Tan, D. X., Reiter, R. J. (2014) Cyclic 3-hydroxymelatonin, a key metabolite enhancing the peroxy radical scavenging activity of melatonin. *RSC Advances*, 4(10): 5220. DOI:10.1039/c3ra44604b.
- Herlinger, A. W., Wenholt, S. L., Long, T. V. (1970) Infrared spectra of amino acids and their metal complexes. II. Geometrical isomerism in bis(amino acidato)copper(II) complexes. *Journal of the American Chemical Society*, 92(22): 6474–6481. DOI:10.1021/ja00725a015.
- Inomata, Y., Shibata, A., Yukawa, Y., Takeuchi, T., Moriwaki, T. (1988) The metal complexes of aminoacids and their N-substituted derivatives—VII. The i.r. spectra and normal coordinate analyses of bivalent metal complexes with N-methylglycine and N-phenylglycine. *Spectrochimica Acta Part A: Molecular Spectroscopy*, 44(1): 97–107. DOI:10.1016/0584-8539(88)80265-4.
- Kruck, T. P. A., Sarkar, B. (1973) Structure of the Species in the Copper (II)–L-Histidine System. *Canadian Journal of Chemistry*, 51(21): 3563–3571. DOI:10.1139/v73-531.
- Kumar, S., Rai, A. K., Rai, S. B., Rai, D. K. (2010) Infrared and Raman spectra of Histidine: an ab initio DFT calculations of Histidine molecule and its different protonated forms. *Indian Journal of Physics*, 84(5): 563–573 DOI:10.1007/s12648-010-0039-6.
- Liu, H., Qu, Y., & Wang, X. (2018). Amyloid β -targeted metal complexes for potential applications in Alzheimer's disease. *Future medicinal chemistry*, 10(6), 679–701. DOI: [10.4155/fmc-2017-0248](https://doi.org/10.4155/fmc-2017-0248)

- Martí, I., Ferrer, A., Escorihuela, J., Burguete, M. I., Luis, S. V. (2012) Copper(II) complexes of bis(amino amide) ligands: effect of changes in the amino acid residue. *Dalton Transactions*, 41(22):6764. DOI:10.1039/c2dt12459a.
- Nakamoto, K. (2009) *Infrared and Raman Spectra of Inorganic and Coordination Compounds: Part A, Theory and Applications in Inorganic Chemistry*. 6th ed. New Jersey: John Wiley and Sons, Inc.
- Nakamoto, K. (2009) *Infrared and Raman Spectra of Inorganic and Coordination Compounds: Part B, Applications in Coordination, Organometallic and Bioinorganic Chemistry*. 6th ed. New Jersey: John Wiley and Sons, Inc.
- Romero-Canelon, I., & Sadler, P. J. (2013). Next-generation metal anticancer complexes: multitargeting via redox modulation. *Inorganic chemistry*, 52(21), 12276-12291. DOI: [10.1021/ic400835n](https://doi.org/10.1021/ic400835n)
- Scheiber, I., Dringen, R., Mercer, J. F. B. (2013) *Copper: Effects of Deficiency and Overload*. In: Interrelations Between Essential Metal Ions and Human Diseases, Dordrecht: Springer Netherlands, pp: 359–387. DOI:10.1007/978-94-007-7500-8_11.
- Shimazaki, Y., Yajima, T., Takani, M., Yamauchi, O. (2009) Metal complexes involving indole rings: Structures and effects of metal–indole interactions. *Coordination Chemistry Reviews*, 253(3-4): 479–492. DOI:10.1016/j.ccr.2008.04.012.
- Simunkova, M., Alwasel, S. H., Alhazza, I. M., Jomova, K., Kollar, V., Rusko, M., & Valko, M. (2019). Management of oxidative stress and other pathologies in Alzheimer's disease. *Archives of toxicology*, 93(9), 2491-2513. DOI: [10.1007/s00204-019-02538-y](https://doi.org/10.1007/s00204-019-02538-y)
- Van Rijt, S. H., & Sadler, P. J. (2009). Current applications and future potential for bioinorganic chemistry in the development of anticancer drugs. *Drug discovery today*, 14(23-24), 1089-1097. DOI: [10.1016/j.drudis.2009.09.003](https://doi.org/10.1016/j.drudis.2009.09.003)
- Vest, K. E., Hashemi, H. F., Cobine, P. A. (2012) The Copper Metallome in Eukaryotic Cells. *Met Ions Life Sci.*, 12: 451–478 DOI:10.1007/978-94-007-5561-1_13.
- Waggoner, D. J., Bartnikas, T. B., Gitlin J. D. (1999), The Role of Copper in Neurodegenerative Disease. *Neurobiology of Disease*, 6(4): 221-230. DOI: 10.1006/nbdi.1999.0250
- Wagner C. C., Baran E. J. (2004) Spectroscopic and magnetic behavior of the Copper(II) complex of L-tryptophan. *Acta Farm. Bonaerense*, 23(3): 339-42. ISSN 0326-2383
- Wang, J., Wang, X., He, Y., Jia, L., Yang, C. S., Reiter, R. J., Zhang, J. (2019) Antioxidant and Pro Oxidant Activities of Melatonin in the Presence of Copper and Polyphenols In Vitro and In Vivo. *Cells*, 8(8):903. DOI:10.3390/cells8080903.
- Zuo, L., Zhou, T., Pannell, B. K., Ziegler, A. C., & Best, T. M. (2015). Biological and physiological role of reactive oxygen species—the good, the bad and the ugly. *Acta physiologica*, 214(3), 329-348. DOI: [10.1111/apha.12515](https://doi.org/10.1111/apha.12515)

Summary/Sažetak

Ispitana je reakcija bezvodnog bakar(II) hlorida u metanolnom rastvoru sa alaninom, glicinom, histidinom, triptofanom kao i reakcija s melatoninom. Produkti reakcija ispitani su FTIR spektroskopijom kako bi se izvršila distinkcija produkata reakcije u odnosu na, u literaturi, prethodno opisane komplekse istih aminokiselina sa Cu(II) kao što su *cis*- i *trans*-bis(glicinato)bakar(II). Od velikog interesa bila je reakcija CuCl₂ s melatoninom u ovim uvjetima. FTIR karakterizacija sintetiziranih kompleksa pokazala je da ovi reakcioni uslovi rezultiraju nastankom kompleksa Cu(II) s glicinom i s alaninom koji su složenije građe od bis(glicinato)- i bis(alaninato)bakar(II) kompleksa. Kompleks priređen s melatoninom pokazuje interesantne trake u spektralnom području za koje je odgovoran atom azota iz amido grupe melatonina. Posljednje opažanje je vrlo značajno jer je poznat vrlo mali broj metalnih kompleksa melatonina koji su dobro okarakterizirani, dok je sinteza većine njih eksperimentalno zahtjevan proces uz simultanu kontrolu više faktora. Ispitivanjem antioksidativnog kapaciteta sintetiziranih kompleksa bakra dobijene su vrijednosti CEAC u rasponu od 121.6 - 734.4 μM. Najniže vrijednosti antioksidativog kapaciteta su zabilježene kod kompleksa bakra sa triptofanom, dok su najviše vrijednosti zabilježene kod kompleksa bakra sa alaninom. Uočen je i visok antioksidativni kapacitet kompleksa bakra sa melatoninom (673.8 μM).